

From:

To:

Date: 2/2/2022 5:53:53 AM

Subject:

Attachments: .thmx

®
(GEPACEF)

1

1,0

AT J01D D12.

®

® *in vitro*

®

Staphylococcus aureus (), Staphylococcus epidermidis, Streptococcus pneumoniae (), Streptococcus agalactiae (- B), Streptococcus faecalis (), -

Escherichia coli, Klebsiella, Enterobacter, Citrobacter, Haemophilus influenzae, Proteus mirabilis, Proteus vulgaris, Morganella morganii (- Proteus morganii), Providencia rettgeri (- Proteus rettgeri), Providenci, Serratia (S. Marcescens), Salmonella, Shigella, Pseudomonas aeruginosa Pseudomonas, Acinetobacter calcoaceticus, Neisseria gonorrhoeae (,), Neisseria meningitidis, Bordetella pertussis, Yersinia enterocolitica.

(*Peptococcus*, *Peptostreptococcus*, *Veillonella*);
Clostridium, *Eubacterium*, *Lactobacillus*);
Fusobacterium, *Bacteroides fragilis* *Bacteroides*).

15- . 1 1, 2, 3 4 (/) 1.

,	0*	30	1	2	4	8	12
1	153	114	73	38	16	4	0,5
2	252	153	114	70	32	8	2
3	340	210	142	89	41	9	2
4	506	325	251	161	71	19	6
1 ,	32**	52	65	57	33	7	1
2 ,	40**	69	93	97	58	14	4

* , ()

** , 15

®

2

() ,

12

20–30 %.

1

6000

1

15

1000 / 2

2200 /
2

®

®

(R)

5

,
,

(. . « »).

1

2-4

()

Clostridium difficile [*Clostridium difficile* associated diarrhea] (CDAD),

C. difficile.

C. difficile

A B,

CDAD.

C. difficile

CDAD

CDAD

2

®

34

10

12-16, 8²⁻⁴, (12¹² 8),
12-16, 8³, (12⁸ 500),
,

®

,
3,5 /100, 4.
18 /
3,5 /100, 4.
®, 2
(2-4)

, 2
8-12). 300 /
12⁵⁰ (200¹ «»).
,

(8)

12¹ .
12¹ .
®, (1) 20-100

		I ,	,	II , , 2 %	,	,
1	, /	,	,	,	,	*
	250	2,6	0,9		4	
	333	1,8	0,6		3	

24 2-8° ,

1

. 2

2 8°.

®

®

®

1,0

, 10

• « . ».

,01032, . , . ,139.

®
(GEPACEF)

1

1,0

III AT J01D D12.

®

in vitro

®

Staphylococcus aureus (), *Staphylococcus epidermidis*, *Streptococcus pneumoniae* (), *Streptococcus agalactiae* (- *Diplococcus pneumoniae*), *Streptococcus pyogenes* (-), *Streptococcus faecalis* (), -

Escherichia coli, *Klebsiella*, *Enterobacter*, *Citrobacter*, *Haemophilus influenzae*, *Proteus mirabilis*, *Proteus vulgaris*, *Morganella morganii* (*Proteus morganii*), *Providencia rettgeri* (*Proteus rettgeri*), *Providencia*, *Serratia* (*S. Marcescens*), *Salmonella*, *Shigella*, *Pseudomonas aeruginosa* *Pseudomonas*, *Acinetobacter calcoaceticus*, *Neisseria gonorrhoeae* (,), *Neisseria meningitidis*, *Bordetella pertussis*, *Yersinia enterocolitica*.

Bacteroides fragilis ((*Peptococcus*, *Peptostreptococcus*, *Velionella*); *Clostridium*, *Eubacterium*, *Lactobacillus*); (*Fusobacterium*, *Bacteroides*).

»).

/

2-4

()

[*Clostridium difficile* associated diarrhea] (CDAD),

C. difficile

A B,

CDAD.

C. difficile

Clostridium difficile

C. difficile.

CDAD

CDAD

2

®

34

, 10

^⑧ 12-16 , 8 , 3 (12 12 8), 500 ().

2

®
12 (. « 50 200 1 ; 2 (8-12).
300 / ,
. .
(8) 12 .

		I /	, ,	II /	, ,	2 %	*
1	250		2,6		0,9		4

*

,

,

$$\begin{array}{c} \text{5} \\ \hline 50 /); 5 \% & 0,2 \% \\ (2 \quad 50 /); & -R (2 \quad 50 /); \\ \textcircled{R} & \end{array} \quad \begin{array}{c} (2-8^\circ) \\ (300 /); 5 \% \\ (300 /); 0,9 \% \\ -R (2 \quad 50 /); \\ (300 /) \end{array} \quad \begin{array}{c} (2 \quad 50 /); 5 \% \\ (2 \quad 300 /); \\ (300 /) \end{array} \quad \begin{array}{c} 0,9 \% \\ -M \quad 5 \% \\ (2 \quad /); 0,5 \% \end{array}$$

$$\left(\begin{array}{cc} -20 & -10^\circ \\ 0,2 \% & (300 /) \end{array} \right) \quad 3 \quad : 5 \% \quad (2 /), \quad \left(\begin{array}{cc} (50 /) 5 \% & 0,9 \% \\ 5 & 0,9 \% \end{array} \right) \quad (300 /) \quad , \quad (2 /) 5 \%$$

1

24

2-8° ,

/10); (1/1000 <1/100); (1/10 000 <1/1000); (CIOMS III): (<1/10 000) (1/10); (1/100 <1).

¹ See, for example, the discussion in *Principles of Corporate Finance*, 2d ed., by M. J. Jensen and William H. Meckling (New York: McGraw-Hill, 1976), pp. 11-12.

. 2

2 8°

®

®,

1,0

, 10

«

».

, 01032, . . . , . . . , 139.